




## **CERTIFICATE**

I, Shigeaki YAMAZAKI, a citizen of Japan, residing at THREE F MINAMI AOYAMA BLDG. 7F, 6-11-1, MINAMI-AOYAMA, MINATO-KU, TOKYO, JAPAN, hereby certify that I am conversant with the English and Japanese language, and I further certify that to the best of my knowledge and belief the foregoing is a true and correct English translation of the Japanese Patent Application No.167439/1999, attached hereto.

Signed this 24<sup>th</sup> day of December, 2004

  
\_\_\_\_\_  
Shigeaki YAMAZAKI



[Name of The Document]	Drawing	1 copy
[Name of The Document]	Abstract	1 copy

[Document] SPECIFICATION

[Title of Invention] Thermogenic Genes and Thermogenic Proteins in Plant

[Claims]

[Claim 1] A thermogenic gene SfUCPa derived from skunk cabbage, of which cDNA comprises the base sequence of SEQ ID NO: 1.

[Claim 2] A thermogenic gene SfUCPb derived from skunk cabbage, of which cDNA comprises the base sequence of SEQ ID NO: 3.

[Claim 3] A thermogenic protein SfUCPA expressed from the gene SfUCPa of Claim 1, which comprises the amino acid sequence of SEQ ID NO: 2.

[Claim 4] A Thermogenic protein SfUCPB expressed from the gene SfUCPb of Claim 2, which comprises the amino acid sequence of SEQ ID NO: 4.

[Claim 5] A DNA fragment comprising the base sequence of SEQ ID NO: 1 or a partial sequence thereof.

[Claim 6] A DNA fragment comprising the base sequence of SEQ ID NO 3: or a partial sequence thereof.

[Detailed Description of Invention]

[0001]

[Technical Field of Invention]

The present invention relates to plant thermogenic genes and proteins. More particularly, the invention relates to thermopogenic genes derived from a skunk cabbage (*Symplocarpus foetidus*) and gene products (proteins). Those genes and proteins are useful in breeding of cold-avoidance plants, medical treatment of diabetes mellitus or obesity, or development of novel thermogenic bio-materials.

[0002]

[Prior Art]

Stresses due to low temperatures, droughts and salinity are common harmful environmental factors that terrestrial plants encounter. Among these stresses, it has been considered that cellular injury due to the low temperature is the most important factor which restricts productivity of crops (Levitt, 1980). To

resist the low temperature stress, cold-hardy plants such as wheat or rye have a variety of physiological and metabolic responses which lead to cold acclimation (Sakai and Larcher, 1987; Steponkus, 1984; Thomashow, 1998; Uemura and Steponkus, 1997). In contrast, it is known that some plants including skunk cabbage have a specialized system by which the plants generate heat to avoid freezing (Knutson, 1974; Nagy et al., 1972; Schneider and Buchenen, 1980).

[0003]

The temperature of the flower in the spadix of skunk cabbage, which flowers in early spring, has been known to maintain its temperature at higher than +10°C even when the ambient temperature falls to -15°C (Knutson, 1974). For example, Fig. 1 is a result of thermoscopic analysis for the spadix using infrared camera, in which the plants were placed in the growth chamber and the air temperature was gradually decreased. As clearly seen from Fig. 1, the temperature of the spadix of skunk cabbage is kept at approximately 19°C notwithstanding a fall of the ambient temperature.

[0004]

The temperature is thus maintained by doubling the respiration rate from the level of 12°C to that of sub-zero temperature. It has also been considered that the heat production in thermogenic plant species relates to a cellular metabolism called cyanide-non-sensitive/non-phosphorylating electron-transferring pathway, which is controlled by mitochondrial alternative oxidase (AOX)(Berthold and Siedow, 1993; Ito et al., 1997; McIntosh, 1994, Wangner and Krab, 1995).

[0005]

On the other hand, it has been shown that a mitochondrial protein called an uncoupling protein (UCP) plays an important role in generation of heat in mammals. UCP found in the intima of mitochondria make H<sup>+</sup> flow into the membrane to uncouple aspiration from synthesis of ATP which acts to disperse chemical energy to metabolic heat (Klaus et al., 1991; Klingenberg and Winkler, 1985; Ricquier et al., 1991). In animals, 3 types of UCPs have been found. UCP1 is primarily distributed in brown adipose tissue (Nichollus and Locke, 1984).

UCP2 is found ubiquitously in many tissues (Fleury et al., 1997), and UCP3 is localized specifically in skeletal muscle (Boss et al., 1997).

[0006]

It has been considered that UCPs of mammals, similarly to other carrier proteins of mitochondria, are composed of 6 transmembrane segments, of which the hydrophobic portion is derived from pairing amphipathic  $\alpha$ -helix structure (Liu et al., 1988; Maia et al., 1998). It is also known that the activity of these UCPs decreases depending on purine nucleotides (ATP, GTP, GDP and ADP) attached to the C-terminal region and increases by free fatty acids (Jezek et al., 1998; Lin and Klingenberg, 1982; Katiyar and Shrago, 1989; Rial et al., 1983; Sluse et al., 1998).

[0007]

On the contrary, 2 cDNAs encoding UCP-like proteins of plant origin were isolated from potato (StUCP: Laloi et al., 1997) and from *Arabidopsis* (AtPUMP: Maia et al., 1998). Since the expression of StUCP was mainly detected in the flower and the fruit, it has been postulated that StUCP may concern respiration during flowering and maturation of the fruit together with the AOX activity (Laloi et al., 1997).

[0008]

Potato and *Arabidopsis* have been considered to be non-thermogenic plants. However, the expression of StUCP and AtPUMP induced by low temperature. Therefore, it has been suggested that these genes are involved in the heat production (Laloi et al., 1997; Maia et al., 1998).

In the thermogenic plants such as skunk cabbage, however, UCP-mediated thermoigenic mechanisms have not yet been identified.

[0009]

[Purpose of Invention]

The purpose of the invention of this application is to provide unidentified novel UCP genes derived from a thermogenic plant, skunk cabbage.

The additional purpose of this application is to provide skunk cabbage UCPs which are expression products of the novel genes.

[0010]

[Solution of the Purpose]

The invention provides thermogenic genes derived from skunk cabbage, i.e., gene SfUCPa of which cDNA comprises the base sequence of SEQ ID NO: 1, and gene SfUCPb of which cDNA comprises the base sequence of SEQ ID NO: 3.

[0011]

Moreover, the invention provides thermogenic proteins, i.e., protein SfUCPA expressed from SfUCPa, which comprises the amino acid sequence of SEQ ID NO: 2, and protein SfUCPB expressed from SfUCPb, which comprises the amino acid sequence of SEQ ID NO: 4.

[0012]

In addition, the invention provides cDNA having the base sequence of SEQ ID NO: 1 or a partial sequence thereof, and cDNA having the base sequence of SEQ ID NO: 3 or a partial sequence thereof.

In the gene SfUCPa of the present invention, its cDNA has the base sequence of SEQ ID NO: 1 and encodes the protein SfUCPA having the amino acid sequence of SEQ ID NO: 2, of which the estimated molecular weight is 32.6 kDa. In the gene SfUCPb of the present invention, its cDNA (SEQ ID NO: 3) encodes the protein SfUCPB having the amino acid sequence of SEQ ID NO: 4, of which the estimated molecular weight is 29.0 kDa.

[0013]

Genes SfUCPa and SfUCPb of the invention are derived from skunk cabbage, which are expressed specifically in the spadix when the temperature is low. The results of Northern blotting on the total RNAs extracted from skunk cabbage (Ito et al., 1999), confirmed that the expressions of both genes were detected in the spadices but not in the leaves at room temperature (15°C) as shown on Fig. 2. It was also confirmed that the spadix-specific expression of both genes were induced by cold treatment (4°C for 3 days).

[0014]

The amino acid sequences of the proteins SfUCPA and SfUCPB that are

expressed from the respective genes of the invention have higher homology to the plant UCPs than to the human UCPs (Fig. 3) such that the amino acid sequence of SfUCPA has homology of 79%, 75%, 44%, 48% and 48% to StUCP, AtPUMP, human UCP, UCP2 and UCP3, respectively. SfUCPB has homology of 71%, 66%, 41%, 43% and 44% to StUCP, AtPUMP, human UCP, UCP2 and UCP3, respectively.

[0015]

In addition, SfUCPA and SfUCPB have high sequence homology (88%) to each other though the region corresponding to the amino acid sequence between the 204th Thr and the 238th Val in SfUCPA is completely deleted in SfUCPB as shown on Fig. 3. Moreover, the 265th Leu of SfUCPA is replaced by Pro in SfUCPB.

[0016]

StUCPA has similar structure to that of other mitochondria UCP proteins. SfUCPA has 6 transmembrane domains as shown by the hydrophobic plot in Fig. 4, of which the topology is as shown in Fig. 5. In addition, this SfUCPA has 3 domains that are characteristic of energy transfer proteins in mitochondria (Fig. 3)(Boss et al., 1997; Maia et al., 1998). On the other hand, SfUCPB is lacking in the 3rd domain which is characteristic of energy transfer proteins in mitochondrial (Fig. 3), as well as in the 5th transmembrane domain (Figs. 3 and 6). The topology is located toward the mitochondria matrix at the C-terminal (Fig. 7).

[0017]

Each protein has a purine nucleotide-binding domain (PNBD) at the C-terminal (Figs. 3, 5 and 7), and it is known that in UCP, binding of the purine nucleotide inhibits the uncoupling function in the mitochondria intima. In SfUCPB, however, there is a possibility that it may have escaped the inhibition of the binding of the purine nucleotide because its C-terminal is located toward the mitochondria matrix. Such a topology has not been found in any UCPs from animals or plants.

[0018]



The thermogenic genes SfUCPa and SfUCPb provided by the invention are derived from skunk cabbage and are very useful in, for example, development of low temperature-tolerant plants using a genetic recombination technique. The proteins SfUCPA and SfUCPB that are expression products from the above genes are expected as effective components in remedies of diabetes mellitus, obesity, and the like, based on the uncoupling function to ATP synthesis. Moreover, such thermogenic proteins are also promising novel heat generating bio-materials.

[0019]

The followings are more detailed description for embodiments of the invention.

[0020]

[Embodiments of Invention]

The genes SfUCPa and SfUCPb of the invention can be isolated from the genomic DNA of skunk cabbage using the cDNA (SEQ ID NOS: 1 or 3) or a partial sequence thereof of the invention as a probe. For example, a genome library is prepared from the genomic DNA according to a known method. It may be screened by means of colony or plaque hybridization according to a known method using as a probe an oligonucleotide synthesized based on the base sequence of an optional portion of cDNA. Alternatively, the target genetic region may also be identified by means of in situ hybridization for chromosome.

[0021]

The respective cDNAs of the invention can be cloned, for example, from a cDNA library which is synthesized using a poly(A)<sup>+</sup>RNA of skunk cabbage as a template. In such a case, an oligonucleotide of an optional portion of cDNA provided by the invention is synthesized, which may be used as a probe to carry out screening by means of colony or plaque hybridization according to a known method. Alternatively, oligonucleotides which can hybridize to both ends of the target cDNA fragment are synthesized, which may be used as primers in preparation of cDNA of the invention by the RT-PCR method from mRNA isolated from the cells of skunk cabbage.

[0022]

In general, polymorphism is frequently recognized in the genes of eucaryotic cells. In the invention, accordingly, in addition to cDNAs represented by SEQ ID NOS: 1 and 3, those in which one or several nucleotides are added, deleted and/or replaced by (an)other nucleotide(s) in the above cDNA are included. Similarly, proteins in which one or more amino acids are added, deleted and/or replaced by (an)other amino acid(s) due to change of the above nucleotide are also included in the present invention.

[0023]

In cDNA of the invention, DNA fragments (10bp or more) comprising an optimal part of the base sequences of SEQ ID NOS 1 and 3 are included. In addition, DNA fragments comprising a sense strand or anti-sense strand are also included.

The proteins of the invention, SfUCPA and SfUCPB, may be prepared respectively by a known method, for example, isolation from the spadix of skunk cabbage, preparation by chemical syntheses based on the amino acid sequence provided by the invention, or production by a recombinant DNA technique using cDNA provided by the invention. For example, when the protein is produced by a recombinant DNA technique, RNA is prepared from a vector containing cDNA of the invention by *in vitro* transcription, and this is used as a template for *in vitro* translation to yield the protein. Alternatively, the translational region of cDNA is incorporated into an appropriate expression vector according to a known method, and the resulting recombinant vector is introduced into *Escherichia coli*, *Bacillus subtilis*, yeast, animal or plant cells. The resulting transformants can be used in expression of the proteins in a large quantity.

[0024]

In the case of the proteins of the invention being produced by *in vitro* translation, the translation region of cDNA of the invention may be incorporated into a vector containing RNA polymerase promotor, and then added to an *in vitro* translation system such as a rabbit reticulocyte lysate or wheat germ extract

containing an RNA polymerase corresponding to the promotor. The RNA polymerase promotor is exemplified by T7, T3, SP6, and similar promotors.

[0025]

In the case of the proteins of the invention being expressed in a microorganism such as *Escherichia coli*, the translation region of cDNA is incorporated into an expression vector containing an origin replicable in microorganisms, promoter, ribosome binding site, cDNA cloning site, terminator, and the like, to construct a recombinant expression vector, which is then introduced into a host cell and incubated. In this operation, an initiation codon and a stop codon may be added to the front and tail of an optional translation region to obtain a protein fragment containing the optional region. Alternatively, the desired protein may be expressed as a fusion protein with another protein, which may be cleaved with a suitable protease to yield the desired protein. The expression vectors for *Escherichia coli* are exemplified by pUC series, pBluescript II, pET expression system, pGEX expression system, and the like.

[0026]

In the case of the proteins of the invention being expressed in eucaryotic cells, the translation region of cDNA of the invention is incorporated into an expression vector for eucaryotic cells containing a promoter, splicing region, poly(A) additional site, and the like, and introduced into the eucaryotic cells. The expression vector is exemplified by pKA1, pCDM8, pSVK3, pMSG, pSVL, pBK-CMV, pBK-RSV, EVB-vector, pRS, pYES2, and the like. The eucaryotic cells, many include mammal cultured cells such as monkey renal cell COS7, Chinese hamster ovarian cell CHO, etc., budding yeast, fission yeast, silkworm cell, *Xenopus* egg cell, and the like are commonly used, but not limited thereto. In order to introduce the expression vector into eucaryotic cells, a known method such as electroporation, calcium phosphate method, liposome method, DEAE dextran method, and the like can be utilized.

[0027]

After expression of the proteins in procaryotic cells or eucaryotic cells

according to the aforementioned method, the desired proteins are isolated and purified from the culture in the known combined procedures for separation. For example, treatment with a denaturant (e.g., urea) or surface activator, ultrasonication, digestion with enzymes, salting-out or solvent precipitation, dialysis, centrifugation, ultrafiltration, gel filtration, SDS-PAGE, isoelectric focusing, ion-exchange chromatography, hydrophobic chromatography, affinity chromatography, reverse phase chromatography, and the like are involved.

[0028]

The proteins of the invention, SfUCPA and SfUCPB, also include peptide fragments (5 amino acids or more) involving the optional partial amino acid sequences of SEQ ID NOS: 2 and 4. In addition, the proteins of the invention also include fusion proteins with other optional proteins.

[0029]

#### [Examples]

The following examples serve to illustrate the invention of this application specifically in more detail, but are not intended to limit the scope of the invention.

#### Example 1: Cloning of cDNA

The total RNA was extracted from the spadix of skunk cabbage (*Symplocarpus foetidus*) and the complete RNA was determined on 1.0% agarose gel electrophoresis (Ito et al., 1999). Using a mRNA isolation kit (Pharmacia), a clone associated with the UCP gene family was isolated from the purified poly(A)+RNA by RT-PCR. The first strand cDNA was prepared by annealing 20 pmol of cDNA primed primer (5'-TTTTTTTTTTTTTTTTTTTTTTTTTTT-3') into poly(A)+RNA (0.1µg), followed by extension with 10 units of reverse transcriptase (New England Biolab) at 37°C for 30 minutes in 20µl of 1×RT buffer containing 10mM 1,4-dithiothreitol and 0.5mM dNTP. The composition of the reaction solution is as follows.

[0030]

- 10mM Tris-HCl (pH 8.0);
- 50mM KCl;

- 1.5mM MgCl<sub>2</sub>;
- 4mM dNTP;
- 0.2 unit of EX Taq polymerase (Takara); and
- 10pmol of two degenerate primers corresponding to the conserved amino acid sequence of the UCP family:

ZF1 (5'-CCIYTIGAYACIGCIAAR-3')

ZR1 (5'-ACWTTCCAISYICCIAWIC-3')

PCR cycle was carried out as follows.

[0031]

(94°C: 0.5 minute; 50°C: 1 minute; 72°C: 1 minute)×35

Among the PCR products obtained in the above method, the amino acid sequence estimated from the sequence of about 0.8kb cDNA fragment had very high homology to one of the reading frame sequences of the UCP gene family. This fragment, accordingly, was cloned into T-vector (clone p2-1) and used as a probe for library screening.

[0032]

cDNA (5μg) prepared from the spadix was inserted into λgt11 phage according to the known method (Sambrook et al., 1989) to construct a cDNA library. From this library, 8 clones positive to the above-described probes were isolated and sub-cloned into the pBluescript SK plasmid (Stratagene). From these clones, clones pz8-1 and pz8-2 were obtained, which respectively contained the full length SfUCPa cDNA and SfUCPb cDNA.

[0033]

The insert in each clone was sequenced with an auto-sequencer ABI373A using the BcaBest sequencing kit (Takara) and T3, T7 and gene-specific primers. The sequence data were analyzed by means of the GENETYX-Homology Software System version 2.2.0 (Software Development).

cDNA of SfUCPa had the 1,525bp base sequence of SEQ ID NO: 1, and cDNA of SfUCPb had the 2,991bp base sequence of SEQ ID NO: 3. An estimated polyadenylated signal (aataaa) was found upstream of 236bp from the poly(A)

sequence in cDNA of SfUCPa, while in cDNA of SfUCPb two polyadenylated sites were recognized at the positions of 1,171bp and 1,243bp. It is noteworthy that cDNA of SfUCPb has a longer 3'-untranslation region than that of SfUCPa.

[0034]

cDNA of SfUCPa had an open reading frame (ORF) encoding 303 amino acids as shown in SEQ ID NO: 1, and this ORF was found to encode the protein SfUCPA of the estimated molecular weight 32.6kDa having the amino acid sequence of SEQ ID NO: 2. On the other hand, cDNA of SfUCPb had an ORF corresponding to 268 amino acids as shown in SEQ ID NO: 3, and found to encode the protein SfUCPB of the estimated molecular weight 29.0kDa.

[0035]

Moreover, it was confirmed from the results of Southern blot analysis that the genome of skunk cabbage contains multiple copies of SfUCPa gene and a single copy of SfUCPb (data not shown).

Example 2: *In vitro* translation of cDNA

The plasmid clones pz8-1 and pz8-2 obtained in Example 1 were linearized, on which a sense- or anti-sense RNA was transcribed with T7 RNA polymerase or T3 RNA polymerase according to the protocol of MAXICRIPT transcription kit (Ambion). An equal amount of RNA (4μg) was provided for *in vitro* translation reaction using a wheat germ extract (Promega) in the presence of <sup>35</sup>S-methionine (Amersham). The translation product was analyzed by SDS-PAGE. The gel was fixed, incubated in Amplify (Amersham), then dried, and fluorometrically analyzed.

[0036]

As a result, it was confirmed that, as shown in Fig. 8, the initiation codon and the stop codon of cDNA isolated in Example 1 functioned successfully since a protein having an expected molecular weight was produced from any of cDNAs only when the sense RNA was used as a template.

[0037]

[Effects of Invention]

As described previously, this application provides novel thermogenic genes SfUCPa and SfUCPb as well as their gene products, i.e., thermogenic proteins SfUCPA and SfUCPB, derived from skunk cabbage (*Symplocarpus foetidus*), and cDNAs used for gene engineering mass production of these proteins. These genes and proteins allow development of low temperature-tolerant plants, development of drugs or methods for treatment of diabetes mellitus or obesity, or development of novel heat generating materials from plants.

[0038]

#### SEQUENCE LISTING

<110> Japan Science and Technology Corporation

<120> Thermogenesis Genes and Thermogenesis Proteins in Plant

<160> 4

<170> PatentIn Ver. 2.0

<210> 1

<211> 1525

<212> DNA

<213> *Symplocarpus foetidus*

<220>

<221> CDS

<222> (280).. (1188)

<221> poly A site

<221> (1271).. (1276)

<300>

<301> Ito, K.

<302> Isolation of two distinct cold-inducible cDNAs encoding plant uncoupling proteins from the spadix of skunk cabbage (*Symplocarpus foetidus*)

<303> Plant Sci.

<304> 149

<305>

<306> 167-173

<307> 1999

<308> GenBank AB024733

<309> 2000-02-25

<400> 1

gaggattcgc agaagaaagg ccagaacccg attccttccc gtcttcttct ccttccgccc 60  
aatgcagtt tttcgcagcg gcgtcatcat caagaccctc cgcctttccg cgccaaacgc 120  
cttccacccc cacccaatcg ccctccgttt cccgaaatat tcctcttccc tcctcccttt 180  
tcttctctac ataaacccta accaccccat cctctcctcc cgcttccgac caccctgcat 240  
tctactggga gcccatitga tcgaggtttc ccggcgagg atg ggc gat cac ggc 294

Met Gly Asp His Gly

1

5

ccg agg acc gag atc tgc ttt gcc ggc agt tgc cga gca gca ttc gcc 342  
Pro Arg Thr Glu Ile Ser Phe Ala Gly Ser Ser Arg Ala Ala Phe Ala

10

15

20

gct tgc ttc gcc gag ctt tgc acg att ccg ttg gac act gct aaa gtt 390  
Ala Cys Phe Ala Glu Leu Cys Thr Ile Pro Leu Asp Thr Ala Lys Val

25

30

35

agg ctt caa ctc caa aag aaa gca gta aca ggt gat gtg gtg gct ttg 438  
Arg Leu Gln Leu Gln Lys Lys Ala Val Thr Gly Asp Val Val Ala Leu

40

45

50

cca aaa tac agg gga atg ttg ggc act gtt gcc act att gcc agg gag 486  
Pro Lys Tyr Arg Gly Met Leu Gly Thr Val Ala Thr Ile Ala Arg Glu

55

60

65

gaa ggt ttg tgc gca ctc tgg aaa gga att gta ccc ggt ttg cat cgt 534  
Glu Gly Leu Ser Ala Leu Trp Lys Gly Ile Val Pro Gly Leu His Arg

70

75

80

85

caa tgc ctc ttt gga ggg cta cga att ggg ttg tat gaa cca gtt aag 582  
Gln Cys Leu Phe Gly Gly Leu Arg Ile Gly Leu Tyr Glu Pro Val Lys

90

95

100

tcc ttt tat gtt gga gat aac ttt gtt gga gat att cct tta tcc aag 630



Ser Phe Tyr Val Gly Asp Asn Phe Val Gly Asp Ile Pro Leu Ser Lys	
105 110 115	
aaa ata ctt gct ggg ctt aca aca ggt gca tta gca att ata gtg gca	678
Lys Ile Leu Ala Gly Leu Thr Thr Gly Ala Leu Ala Ile Ile Val Ala	
120 125 130	
aat ccc act gac ctt gtt aaa gtt cga ctt caa tct gaa ggt aaa ctc	726
Asn Pro Thr Asp Leu Val Lys Val Arg Leu Gln Ser Glu Gly Lys Leu	
135 140 145	
ccc cct ggg gta ccg aga cgt tat tca ggg gcg cta aat gct tat tca	774
Pro Pro Gly Val Pro Arg Arg Tyr Ser Gly Ala Leu Asn Ala Tyr Ser	
150 155 160 165	
acc ata gtc aaa aag gaa gga ctt ggt gct ctg tgg act ggg ctt ggt	822
Thr Ile Val Lys Lys Glu Gly Leu Gly Ala Leu Trp Thr Gly Leu Gly	
170 175 180	
cct aat att gcc cgc aat gct att ata aat gct gct gaa ttg gcc agt	870
Pro Asn Ile Ala Arg Asn Ala Ile Ile Asn Ala Ala Glu Leu Ala Ser	
185 190 195	
tat gat caa gtg aaa cag aca atc tta aaa tta cca gga ttc tca gat	918
Tyr Asp Gln Val Lys Gln Thr Ile Leu Lys Leu Pro Gly Phe Ser Asp	
200 205 210	
aat att ttt act cat att tta gcc ggt ctg ggg gca ggt ttt ttt gcc	966
Asn Ile Phe Thr His Ile Leu Ala Gly Leu Gly Ala Gly Phe Phe Ala	
215 220 225	
gtc tgt atc ggt tct cct gtt gat gtg atg aag tct aga atg atg gga	1014
Val Cys Ile Gly Ser Pro Val Asp Val Met Lys Ser Arg Met Met Gly	
230 235 240 245	
gat tca gcc tac aaa agc aca ttt gat tgt ttc atc aag aca ttg aaa	1062
Asp Ser Ala Tyr Lys Ser Thr Phe Asp Cys Phe Ile Lys Thr Leu Lys	
250 255 260	
aat gat ggg ctt ctt gct ttt tac aag ggg ttt atc cca aac ttt ggt	1110

Asn Asp Gly Leu Leu Ala Phe Tyr Lys Gly Phe Ile Pro Asn Phe Gly  
                   265                                  270                                  275  
 cgg tta gga tcg tgg aat gtg atc atg ttt ttg acc ttg gag cag gtc 1158  
 Arg Leu Gly Ser Trp Asn Val Ile Met Phe Leu Thr Leu Glu Gln Val  
                   280                                  285                                  290  
 aag aag ttt ttc atc aaa gag gtg cca aat taatacattg aactcggata 1208  
 Lys Lys Phe Phe Ile Lys Glu Val Pro Asn  
                   295                                  300  
 ggagtagaaa gaaaggggtt ttgtggaatt ttctctaccg gtgtggatcc tggcgagaga 1268  
 caaataaatc ttctgactg ctccagatgtg tacctttttt atgaatgggt cttttcttat 1328  
 agaggacaga gaaaagaaaa aaaaaatcat tgtcatttac tctttttccc catttctgct 1388  
 gctaattctg gtaggagaag aaaagtctta cattgagtga taacgttggt ctctgcatcc 1448  
 attatttttc agagatacta ttgacacat gaaaagtaat gcacatcagg ttttttttaa 1508  
 aaaaaaaaaa aaaaaaa 1525

<210> 2

<211> 303

<212> PRT

<213> *Symplocarpus foetidus*

<400> 2

Met Gly Asp His Gly Pro Arg Thr Glu Ile Ser Phe Ala Gly Ser Ser  
   1                  5                                  10                                  15  
 Arg Ala Ala Phe Ala Ala Cys Phe Ala Glu Leu Cys Thr Ile Pro Leu  
                   20                                  25                                  30  
 Asp Thr Ala Lys Val Arg Leu Gln Leu Gln Lys Lys Ala Val Thr Gly  
                   35                                  40                                  45  
 Asp Val Val Ala Leu Pro Lys Tyr Arg Gly Met Leu Gly Thr Val Ala  
                   50                                  55                                  60  
 Thr Ile Ala Arg Glu Glu Gly Leu Ser Ala Leu Trp Lys Gly Ile Val  
   65                  70                                  75                                  80

Pro Gly Leu His Arg Gln Cys Leu Phe Gly Gly Leu Arg Ile Gly Leu			
	85	90	95
Tyr Glu Pro Val Lys Ser Phe Tyr Val Gly Asp Asn Phe Val Gly Asp			
	100	105	110
Ile Pro Leu Ser Lys Lys Ile Leu Ala Gly Leu Thr Thr Gly Ala Leu			
	115	120	125
Ala Ile Ile Val Ala Asn Pro Thr Asp Leu Val Lys Val Arg Leu Gln			
	130	135	140
Ser Glu Gly Lys Leu Pro Pro Gly Val Pro Arg Arg Tyr Ser Gly Ala			
145	150	155	160
Leu Asn Ala Tyr Ser Thr Ile Val Lys Lys Glu Gly Leu Gly Ala Leu			
	165	170	175
Trp Thr Gly Leu Gly Pro Asn Ile Ala Arg Asn Ala Ile Ile Asn Ala			
	180	185	190
Ala Glu Leu Ala Ser Tyr Asp Gln Val Lys Gln Thr Ile Leu Lys Leu			
	195	200	205
Pro Gly Phe Ser Asp Asn Ile Phe Thr His Ile Leu Ala Gly Leu Gly			
	210	215	220
Ala Gly Phe Phe Ala Val Cys Ile Gly Ser Pro Val Asp Val Met Lys			
225	230	235	240
Ser Arg Met Met Gly Asp Ser Ala Tyr Lys Ser Thr Phe Asp Cys Phe			
	245	250	255
Ile Lys Thr Leu Lys Asn Asp Gly Leu Leu Ala Phe Tyr Lys Gly Phe			
	260	265	270
Ile Pro Asn Phe Gly Arg Leu Gly Ser Trp Asn Val Ile Met Phe Leu			
	275	280	285
Thr Leu Glu Gln Val Lys Lys Phe Phe Ile Lys Glu Val Pro Asn			
	290	295	300

<210> 3

<211> 2991  
 <212> DNA  
 <213> *Symplocarpus foetidus*  
 <220>  
 <221> CDS  
 <222> (286).. (1089)  
 <221> poly A site  
 <222> (1171) (1176)  
 <221> poly A site  
 <222> (1243) (1248)  
 <300>  
 <301> Ito, K.  
 <302> Isolation of two distinct cold-inducible cDNAs encoding plant uncoupling  
 proteins from the spadix of skunk cabbage (*Symplocarpus foetidus*)  
 <303> Plant Sci.  
 <304> 149  
 <305>  
 <306> 167-173  
 <307> 1999  
 <308> GenBank AB024734  
 <309> 2000-02-25  
 <400> 3

tgggtggtgac gagtgacgag gattcgcaga agaaaggcca gaacccgatt ccttcccgtc 60  
 ttctttctcct tccgcccaat tgcagttttt cgcagcgggt catcatcaag accctccgcc 120  
 tttccgcgcc aaacgccttc cacccaatcc ctccgtttcc cgaaatattc cccttccttc 180  
 ccttttcttc tctacataaa ccctaaccac ccccatcctc tcttcccgct tccgaccacc 240  
 ctgcattcta ctgggatccc atttgatcga cgtttcccg gagg atg ggc gat cac 297

Met Gly Asp His

1

ggc ccg agg acc gag atc tcg ttt gcc ggc agt tcg cga gca gca ttc 345

Gly	Pro	Arg	Thr	Glu	Ile	Ser	Phe	Ala	Gly	Ser	Ser	Arg	Ala	Ala	Phe	
5					10					15					20	
gcc	gct	tgc	ttc	gcc	gag	ctc	tgt	acg	att	ccg	ttg	gac	act	gct	aaa	393
Ala	Ala	Cys	Phe	Ala	Glu	Leu	Cys	Thr	Ile	Pro	Leu	Asp	Thr	Ala	Lys	
				25					30					35		
ggt	agg	ctt	cag	ctc	caa	aag	aaa	gca	gta	aca	ggg	gat	gtg	gtg	gct	441
Val	Arg	Leu	Gln	Leu	Gln	Lys	Lys	Ala	Val	Thr	Gly	Asp	Val	Val	Ala	
				40					45					50		
ttg	cca	aaa	tac	agg	gga	atg	ttg	ggc	act	ggt	gcc	act	att	gcc	agg	489
Leu	Pro	Lys	Tyr	Arg	Gly	Met	Leu	Gly	Thr	Val	Ala	Thr	Ile	Ala	Arg	
		55					60						65			
gag	gaa	ggg	ttg	tgc	gca	ctc	tgg	aaa	gga	att	gta	ccc	ggg	ttg	cat	537
Glu	Glu	Gly	Leu	Ser	Ala	Leu	Trp	Lys	Gly	Ile	Val	Pro	Gly	Leu	His	
	70					75					80					
cgt	caa	tgc	ctc	ttt	gga	ggg	cta	cga	att	ggg	ttg	tat	gaa	cca	gtt	585
Arg	Gln	Cys	Leu	Phe	Gly	Gly	Leu	Arg	Ile	Gly	Leu	Tyr	Glu	Pro	Val	
	85				90					95				100		
aag	tcc	ttt	tat	gtt	gga	gat	aac	ttt	gtt	gga	gat	att	cct	tta	tcc	633
Lys	Ser	Phe	Tyr	Val	Gly	Asp	Asn	Phe	Val	Gly	Asp	Ile	Pro	Leu	Ser	
				105					110					115		
aag	aaa	ata	ctt	gct	ggg	ctt	aca	aca	ggg	gca	tta	gca	att	ata	gtg	681
Lys	Lys	Ile	Leu	Ala	Gly	Leu	Thr	Thr	Gly	Ala	Leu	Ala	Ile	Ile	Val	
			120				125					130				
gca	aat	ccg	act	gac	ctt	gtt	aaa	gtt	cga	ctt	caa	tct	gaa	ggg	aaa	729
Ala	Asn	Pro	Thr	Asp	Leu	Val	Lys	Val	Arg	Leu	Gln	Ser	Glu	Gly	Lys	
		135					140					145				
ctc	ccc	cct	ggg	gta	cca	aga	cgt	tat	tca	ggg	gcg	cta	aat	gct	tat	777
Leu	Pro	Pro	Gly	Val	Pro	Arg	Arg	Tyr	Ser	Gly	Ala	Leu	Asn	Ala	Tyr	
	150					155					160					
tca	acc	ata	gtc	aaa	aag	gaa	gga	ctt	ggg	gct	ctg	tgg	act	ggg	ctt	825

Ser Thr Ile Val Lys Lys Glu Gly Leu Gly Ala Leu Trp Thr Gly Leu  
 165 170 175 180  
 ggt cct aat att gcc cgc aat gct att ata aat gct gct gaa ttg gcc 873  
 Gly Pro Asn Ile Ala Arg Asn Ala Ile Ile Asn Ala Ala Glu Leu Ala  
 185 190 195  
 agt tat gat caa gtg aaa cag atg aag tct aga atg atg gga gat tca 921  
 Ser Tyr Asp Gln Val Lys Gln Met Lys Ser Arg Met Met Gly Asp Ser  
 200 205 210  
 gcc tac aaa agc aca ttt gat tgt ttc atc aag acg ttg aaa aat gat 969  
 Ala Tyr Lys Ser Thr Phe Asp Cys Phe Ile Lys Thr Leu Lys Asn Asp  
 215 220 225  
 ggg cct ctt gct ttt tac aag ggg ttt atc cca aac ttt ggt cgg tta 1017  
 Gly Pro Leu Ala Phe Tyr Lys Gly Phe Ile Pro Asn Phe Gly Arg Leu  
 230 235 240  
 gga tcg tgg aat gtg atc atg ttt ttg acc ttg gag cag gtc aag aag 1065  
 Gly Ser Trp Asn Val Ile Met Phe Leu Thr Leu Glu Gln Val Lys Lys  
 245 250 255 260  
 ttc ttc atc aaa gag gtg cca aat taatacattg aagtcggata ggagtagaaa 1119  
 Phe Phe Ile Lys Glu Val Pro Asn  
 265  
 aaaagggttt ttgtggaatt ttctctaccg gtgtggatcc tggcgagaga gaataaatct 1179  
 tcctgactgc tcagatgttg tacctttttt atgaatgggt cttttcttat agaggacaga 1239  
 gaaaataaaa gaaaaattca ttgtcatgta ctctttttcc ccatttctgc tgagtagcag 1299  
 ctataccaag cagactttgt tgcttggctg ctgctaactt tgtagctgaa gaaaagtctt 1359  
 acattgagtg ataacgttgt tctctgcatc cattattttt cagagttact atttgacaca 1419  
 tgaagagttt tttttttttt tttttttttt aacaggcagc aaatagagga atcgatctca 1479  
 cgactatcct ctttatcat taacaggcat acaacttag ggagagcatg cagggtatat 1539  
 atcaaaatat acccttttat tagacatttt gcgtacacag ttggtcctca aacgactgta 1599  
 tctagcagcc aatttttaga ccacattaag acagagagaa acaagcagaa gaacagggta 1659  
 ccatacatat ataggtaata attaagatga tgaacatagc ataggttcat gatctacttc 1719

ttcttcacgt acacatgatg caccagctga atgggaatct tggtcacat atggcatgaa 1779  
 agtacgtcat gtgcagacgt tatatagtgt tcttcttacc attcagcagc agcaccagag 1839  
 gcatcaaaca ctgggggtctt gacaggggtat gaggggtaca ttgcgatccc acacatgccg 1899  
 tagggagtca cattgcgttt gatttttatg tatccagcct gtccccacct agtgccccat 1959  
 gatttcctta caagccaata atccttccca tcctccgaac catatcctat taccaccaca 2019  
 gcatggtcga tacgttgacc acatgggtcca gcaaatacgc ccgagggtga gtgttggaat 2079  
 ccagcgcccg aagcctcaag agcaacactg acaggttgct ttgcgactgc atactgtagg 2139  
 ctaacctcgt tgtacggaga aacattttca tacgcatcaa tcgagggtgac ttaataaga 2199  
 ttgtctctgc aagttcctcg acgtcctgtg tacgggtaat tttgaaaaa aggagaaggg 2259  
 cgggaagaag cgcgcgtctc tgctcgcgac gggtaattc ttcatatggc cacttcgagc 2319  
 atggcttcgg cagcctccag cttcgttctc actccggcct cccctccacc accacgccgg 2379  
 ttccccgcgc gccttcttt cccagggctg caggaggctc gtgggtggtg gggccgagga 2439  
 agccgcgacg acccccgcgc ccggcgccgg cggagggagc cgcgccgccc cccccaagcc 2499  
 gccaccgatc gggcccaaga ggggtcaaa ggttgatata tagttcttaa tttctttccc 2559  
 gtgtggcttc tggagttaga tttgtttcct ctctctttt tttgtttgt tttttcaatt 2619  
 taaattttat tctcatctgt ggacgacctt ccatcggggt tttcgtccct ctgcgaggtg 2679  
 aagatctccg gaaggaatcc tactggttca acggtgtcgg atcgggtggtg gctgttgatc 2739  
 aggatccggc gactcgatac ccggtcgtgg ttccggtcac caaggtcaac tatgcgaacg 2799  
 tctcgaccaa caactacgca ctggacgaga tcctggaggt gaaatgaggg tcggcgggcg 2859  
 tggtcggctg ggcatgtcac gatgatgtat ttctgcagtt ggtagtgtaa aataccatgt 2919  
 cattcgtgta aaactctttc gttcgccaaa tcctcagttg aaattttaat tcccagccag 2979  
 taaaaaaaa aa 2991

<210> 4

<211> 268

<212> PRT

<213> Symplocarpus foetidus

<400> 4

Met Gly Asp His Gly Pro Arg Thr Glu Ile Ser Phe Ala Gly Ser Ser

1

5

10

15

Arg	Ala	Ala	Phe	Ala	Ala	Cys	Phe	Ala	Glu	Leu	Cys	Thr	Ile	Pro	Leu
		20						25						30	
Asp	Thr	Ala	Lys	Val	Arg	Leu	Gln	Leu	Gln	Lys	Lys	Ala	Val	Thr	Gly
		35					40						45		
Asp	Val	Val	Ala	Leu	Pro	Lys	Tyr	Arg	Gly	Met	Leu	Gly	Thr	Val	Ala
		50					55					60			
Thr	Ile	Ala	Arg	Glu	Glu	Gly	Leu	Ser	Ala	Leu	Trp	Lys	Gly	Ile	Val
	65				70					75				80	
Pro	Gly	Leu	His	Arg	Gln	Cys	Leu	Phe	Gly	Gly	Leu	Arg	Ile	Gly	Leu
			85						90					95	
Tyr	Glu	Pro	Val	Lys	Ser	Phe	Tyr	Val	Gly	Asp	Asn	Phe	Val	Gly	Asp
		100							105					110	
Ile	Pro	Leu	Ser	Lys	Lys	Ile	Leu	Ala	Gly	Leu	Thr	Thr	Gly	Ala	Leu
		115					120						125		
Ala	Ile	Ile	Val	Ala	Asn	Pro	Thr	Asp	Leu	Val	Lys	Val	Arg	Leu	Gln
		130					135						140		
Ser	Glu	Gly	Lys	Leu	Pro	Pro	Gly	Val	Pro	Arg	Arg	Tyr	Ser	Gly	Ala
	145					150				155				160	
Leu	Asn	Ala	Tyr	Ser	Thr	Ile	Val	Lys	Lys	Glu	Gly	Leu	Gly	Ala	Leu
			165						170					175	
Trp	Thr	Gly	Leu	Gly	Pro	Asn	Ile	Ala	Arg	Asn	Ala	Ile	Ile	Asn	Ala
		180							185					190	
Ala	Glu	Leu	Ala	Ser	Tyr	Asp	Gln	Val	Lys	Gln	Met	Lys	Ser	Arg	Met
		195					200							205	
Met	Gly	Asp	Ser	Ala	Tyr	Lys	Ser	Thr	Phe	Asp	Cys	Phe	Ile	Lys	Thr
	210						215							220	
Leu	Lys	Asn	Asp	Gly	Pro	Leu	Ala	Phe	Tyr	Lys	Gly	Phe	Ile	Pro	Asn
	225					230					235			240	
Phe	Gly	Arg	Leu	Gly	Ser	Trp	Asn	Val	Ile	Met	Phe	Leu	Thr	Leu	Glu
			245						250					255	



Gln Val Lys Lys Phe Phe Ile Lys Glu Val Pro Asn

260

265

[Brief Description of Drawings]

[Fig. 1]

A graph showing the change of the temperature of the spadix in skunk cabbage and that of ambient temperature with a lapse of time.

[Fig. 2]

The results of northern blotting, indicating the expression profile of SfUCPa (A) and SfUCPb (B) in the spadix and leaf of skunk cabbage at room temperature (RT) and during cold treatment (4°C for 3 days). The lower figures respectively show the results of ethidium bromide staining of non-decomposed rRNA.

[Fig. 3]

The alignment of amino acid sequences of SfUCPA and SfUCPB, together with potato UCP (StUCP), *Arabidopsis* UCP (AtPUMP) and human UCP. The asterisks (\*) attached under the sequences indicate the same amino acid sequence, and the dot (.) indicates the conservative change in all of the sequences. The boldface indicates the same sequence between SfUCPA and SfUCPB. The gap introduced to optimize the sequence alignment is indicated by a dash (-). The alignment was made using a CLUSTAL W program. The characteristic domains of energy transfer proteins typical of mitochondria are surrounded by a square. The shaded bars (I~VI) above the upper sequence show estimated transmembrane domains.

[Fig. 4]

A hydrophobic plot of SfUCPA. The vertical axis indicates the degree of hydrophobicity and the estimated transmembrane domains are indicated by TM1 to TM6.

[Fig. 5]

A diagrammatic illustration of SfUCPA topology in the mitochondria membrane.

[Fig. 6]

A hydrophobic plot of SfUCPB. The vertical axis indicates the degree of hydrophobicity and the estimated transmembrane domains are indicated by TM1 to TM4 and TM6.

[Fig. 7]

A diagrammatic illustration of SfUCPB topology in the mitochondria membrane.

[Fig. 8]

The results of *in vitro* translation using respective cDNAs of the genes SfUCPa and SfUCPb as templates. (-) indicates a control, S a sense RNA, and AS an antisense RNA. The asterisk (\*) indicates a non-specific product and the empty circle denotes the position of a low molecular translated artificial product synthesized from a small ORF.

#### [References]

- Berthold and Siedow (1993) Plant Physiol. 101, 113-119.
- Boss et al. (1997) FEBS Lett. 408, 39-42.
- Fleury et al. (1997) Nature Genetics 15, 269-272.
- Ito, K. et al. (1999) Plant Sci. 142, 57-65.
- Ito, K. et al. (1994) Nucl. Acids Res. 22, 2036-2041.
- Ito, Y. et al. (1997) Gene 12, 121-129.
- Jezek et al. (1998) Biochem. Biophys. Acta 1365, 319-327.
- Katiyar and Shrago (1989) Nati. Acad. Sci. USA 86, 2559-2562.
- Klaus et al. (1991) Int. J. Biochem. 23, 791-810.
- Klingenberg and Winkler (1985) EMBO J. 4, 3087-3092.
- Knutson (1974) Science 186, 746-747.
- Laloi et al. (1997) Nature 389, 135-136.
- Levitt (1980) Responses of plants to environmental stresses. 2nd edn. New York: Academic Press.
- Lin and Klingenberg (1982) Biochemistry 21, 2950-2956.
- Liu et al. (1988) J. Cell. Biol. 107, 503-509.

- Maia et al. (1998) FEBS Lett. 429, 403-406.
- McIntosh (1994) Plant Physiol. 105, 781-786.
- Nagy et al. (1972) Science 178, 1195-1197.
- Nicholls and Locke (1984) Physiol. Rev. 64, 1-64.
- Rial et al. (1983) Eur. J. Biochem. 137, 197-203.
- Ricquier et al. (1991) FASEB J. 5, 2237-2242.
- Sakai and Larcher (1987) Frost Survival of Plants: Responses and Adaptations to Freezing Stresses. Berlin and New York; Springer-Verlag.
- Sambrook et al. (1989) Molecular Cloning: A Laboratory Manual, 2nd eds. New York; Cold Spring Harbor Laboratory Press.
- Schneider and Buchanan (1980) Amer. J. Bot. 67, 182-193.
- Sluse et al. (1998) FEBS Lett. 433, 237-240.
- Steponkus (1984) Annu. Rev. Plant Physiol. 35, 543-581.
- Thomashow (1998) Plant Physiol. 118, 1-7.
- Uemura and Steponkus (1997) Plant Cold Hardiness: Molecular Biology, Biochemistry, and Physiology (Li and Chen, edn). New York: Plenum Press, pp. 171-179.
- Wagner and Krab (1995) Physiol. Plant, 95, 318-325.

[Document]      ABSTRACT

[Abstract]

[Purpose]      To provide novel UCP genes derived from a thermogenic plant, skunk cabbage, thermogenic proteins which are expression products of the novel genes and cDNAs for mass production of the proteins in genetic engineering.

[Solution]      Thermogenic genes named SfUCPa and SfUCPb derived from skunk cabbage, of which cDNA comprises the base sequence of SEQ ID NO: 1 and 3, respectively, thermogenic proteins, SfUCPA and SfUCPB, expressed from the genes, which comprises the amino acid sequence of SEQ ID NO: 2 and 4, and cDNAs of the genes.

[Representative Drawing]      None